



# Rheology-based sea ice dynamics: from the fluid-like to the state-of-the-art solid-like brittle approach




Niccolò Zanotti<sup>1</sup>, Ivo Pasmans<sup>2</sup>, Alberto Carrassi<sup>1,2</sup>  
<sup>1</sup>Univ. of Bologna, Italy  
<sup>2</sup>Univ. of Reading, UK

## Introduction

The increasing interest in the climate in general and, in particular, in the role that Arctic processes play within it has led to an increasing demand for accurate predictions for sea ice **motion** and **deformation**, i.e. its dynamics. Finding suitable equations to describe the sea ice dynamics has been a long-standing challenge for the sea ice community. The determination of an appropriate constitutive relation, in particular, has been a demanding task. This work reviews the developments in continuum-based sea ice modeling with a particular focus on the formulation of internal stresses, i.e. the **rheology**, going from early fluid-like models to the most recent neXtSIM<sub>DC</sub> model employing the **solid-brittle** rheology.

## A model of sea ice dynamics

- Ice state variable  $J \in \mathcal{J} = \{A, \hat{h}, \dots\}$ , where  $\mathcal{J}$  is **model-dependent**
- Sea ice mean thickness  $\hat{h}$  and concentration  $A$  being the most important
- Sea ice internal stress  $\sigma$ , strain  $\varepsilon$ , strain rate  $\dot{\varepsilon}$
- Equation of motion is depth-integrated




- $F_{ext} = \underbrace{A(\tau_a + \tau_w)}_{\text{air-water stress}} - \underbrace{\rho \hat{h} f_c \hat{k} \times u}_{\text{Coriolis}} - \underbrace{\rho h g \nabla H}_{\text{gravity}}$
- State variables may follow a conservation/transport law

## Fluid-like models

- First efforts to model sea ice rheological behavior: **viscous fluid** model (poor approximation);
- Compact ice deforms sporadically and irreversibly  $\rightarrow$  critical stress states specified by a **yield curve**
- **Plastic** rheologies were developed and coupled with different subcritical behaviors:
  - **elastic** (EP model, Coon et al. 1974)  $\sigma = EK : \varepsilon$
  - **viscous** (VP model, Hibler 1979)  $\sigma = 2\eta\dot{\varepsilon} + [\zeta - \eta]\text{tr}\dot{\varepsilon}\mathbf{I} - \frac{P}{2}\mathbf{I}$
- A flow rule specifies how deformation occurs once a critical stress is reached

## Multifractality of sea ice deformation


- Observed spatial/temporal scaling of sea ice deformation
- **Intermittency** and **heterogeneity** of deformation



## Solid-like models

- Sea ice cover modeled as a progressively damaging medium (with a Mohr-Coulomb criterion). Damage level  $d \leq 1$ ;  $E = E(d)$ ,  $\eta = \eta(d)$ ,  $\lambda = \eta/E$
- **Elasto-brittle (EB)**  $\frac{D\sigma}{Dt} = E(\mathbf{K} : \dot{\varepsilon}) - \frac{d}{1-d}\sigma$
- **Maxwell-Elasto-brittle (MEB)**  $\frac{D\sigma}{Dt} = E(\mathbf{K} : \dot{\varepsilon}) - \frac{\sigma}{\lambda} \left(1 + \lambda \frac{d}{1-d}\right)$
- **Brittle-Bingham-Maxwell (BBM)**  $\frac{D\sigma}{Dt} = E(\mathbf{K} : \dot{\varepsilon}) - \frac{\sigma}{\lambda} \left(1 + \tilde{P} + \lambda \frac{d}{1-d}\right)$

## A timeline



## Summary

- In viscous-fluid models the mechanical parameters ( $\eta$ ,  $E$ ,  $\lambda$ ) are independent of the internal stresses. This resulted in unphysical behaviour.
- In plastic models ice dynamic parameters adjust to keep stresses (sub)critical. However, these models fail to reproduce the observed scaling laws (multifractality).
- Brittle models, in which ice dynamic parameters do not instantaneously relax back after stresses are released, perform better at this point.

## KEY REFERENCES

See full-text for the complete bibliography.

Coon, M.D. et al. (1974). "Modeling the pack ice as an elastic-plastic material". In: *AIDJEX Bulletin* 24, pp. 1–105.

Dansereau, V. et al. (2016). "A Maxwell-elasto-brittle rheology for sea ice modelling". In: *The Cryosphere* 10, pp. 1339–1359.

Girard, L. et al. (2011). "A new modeling framework for sea-ice mechanics based on elasto-brittle rheology". In: *Annals of Glaciology* 52.57, pp. 123–132.


Hibler, W.D. (1979). "A dynamic thermodynamic sea ice model". In: *Journal of Physical Oceanography* 9.4, pp. 815–846.

Ólason, E. et al. (2022). "A New Brittle Rheology and Numerical Framework for Large-Scale Sea-Ice Models". In: *Journal of Advances in Modeling Earth Systems* 14.8.

Richter, T. et al. (2023). "A dynamical core based on a discontinuous Galerkin method for higher-order finite-element sea ice modeling". In: *Geoscientific Model Development* 16.13, pp. 3907–3926.

## ACKNOWLEDGEMENT AND PARTNERS

Schmidt Sciences



## CONTACT INFORMATION



Niccolò Zanotti  
UniBo – DIFA

niccolo.zanotti@studio.unibo.it

<https://github.com/niccolozanotti>